



Pulmonology

Delivery Need By: \_\_\_\_\_ Deliver to:  Patient's Home  Physician's Office  Other \_\_\_\_\_

PATIENT INFORMATION

Patient Name: \_\_\_\_\_  Male
Address: \_\_\_\_\_  Female
City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_
Phone Number: \_\_\_\_\_
Email Address: \_\_\_\_\_
Last Four of Social: \_\_\_\_\_ DOB: \_\_\_\_\_

PRESCRIBER INFORMATION

Prescriber's Name: \_\_\_\_\_
Office Contact Name: \_\_\_\_\_
Address: \_\_\_\_\_
City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_
Phone Number: \_\_\_\_\_ Fax: \_\_\_\_\_
DEA/NPA #: \_\_\_\_\_

INSURANCE - PLEASE FAX COPY OF PRESCRIPTION CARD FRONT & BACK

CLINICAL INFORMATION

Diagnosis: \_\_\_\_\_
ICD-10 Code: \_\_\_\_\_
Height: \_\_\_\_\_ ft \_\_\_\_\_ inches Weight: \_\_\_\_\_
lbs Allergies: \_\_\_\_\_

Has the patient been treated previously for this condition?
 Yes  No
Medications Failed: \_\_\_\_\_
Medications On: \_\_\_\_\_
Other Notes: \_\_\_\_\_

PRESCRIPTION INFORMATION

Table with 5 columns: Medication, Dosage/Strength, Directions, Quantity, Refills. Rows include Adcirca, Bethkis, Cinqair, Dupixent, Kitabis Pak, Pulmozyme.

Physician Signature: \_\_\_\_\_ Date: \_\_\_\_\_



### Pulmonology

Delivery Need By: \_\_\_\_\_ Deliver to:  Patient's Home  Physician's Office  Other \_\_\_\_\_

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Prescriber's Name: \_\_\_\_\_

#### PRESCRIPTION INFORMATION

Medication:	Dosage/Strength:	Directions:	Quantity:	Refills:
Revatio	<input type="checkbox"/> 20 mg tablet <input type="checkbox"/> 10mg/12.5 ml Single-Use Vial <input type="checkbox"/> 10mg/ml when reconstituted	<input type="checkbox"/> Take 20 mg (One Tablet) three times a day <input type="checkbox"/> Other	<input type="checkbox"/> _____ Day supply	
Tobi Podhaler	<input type="checkbox"/> 28mg capsules	<input type="checkbox"/> Inhale contents of four capsules (112 mg) twice daily using Podhaler devices <input type="checkbox"/> Other	<input type="checkbox"/> _____ 28 day multipack <input type="checkbox"/> Other	
Tobramycin	<input type="checkbox"/> 300 mg/5ml ampule	<input type="checkbox"/> Administer one ampule twice daily (12 hours apart) by inhalation in alternating repeated cycles of 28 days on drug, followed by 28 days off drug <input type="checkbox"/> Other:	<input type="checkbox"/> 4 week supply	
Xolair	<input type="checkbox"/> 75 mg/0.5 mL in a single-dose prefilled syringe <input type="checkbox"/> 150 mg/mL solution in a single-dose prefilled syringe <input type="checkbox"/> 150 mg lyophilized powder in a single-dose vial for reconstitution	<input type="checkbox"/> Inject _____ mg every 2 weeks <input type="checkbox"/> Inject _____ mg every 4 weeks <input type="checkbox"/> Other	<input type="checkbox"/> 30 day supply <input type="checkbox"/> 90 day supply <input type="checkbox"/> Other	
<input type="checkbox"/> Patient is interested in patient support programs		<input type="checkbox"/> Ancillary supplies provided for administration		

Physician Signature: \_\_\_\_\_ Date: \_\_\_\_\_